

Revitalizing Therapy

Formula 11 Esophagus-Stomach-Intestines

Problem	Uses	Solution	Composition	Action Mechanisms	Posology
<p>With age, digestive system functionality progressively decreases due to cell damage, which leads to food absorption problems and inefficient digestive process.</p> <p>This deterioration in organ function predisposes to chronic degenerative diseases.</p>	<p>Colitis, Gastritis, malabsorption syndrome, abdominal distension and constipation.</p>	<p>The contribution of cellular cytokines and growth factors in embryonic extracts has a restorative and revitalizing effect at the cellular level, which increases the specific functionality of the organ to be treated.</p> <p>Antioxidant enzymes neutralize free radicals, thereby reducing damage from oxidative stress.</p>	<p>Oral CELLORGANE 11 Each 500-mg enteric coated tablet contains: Opotherapeutic cell extracts: Stomach 15%, Intestine 15%, esophagus 15%, pancreas 15%, embryonic endoderm 10%, Thymus 10%, Placenta 10%; Antioxidant enzyme complex: Superoxide dismutase, glutathione peroxidase, glutathione reductase, glutathione transferase; Maltodextrin 10% and stabilizers.</p> <p>Injectable CELLORGANE 2 Each 750-mg Lyophilized Vial contains: Opotherapeutic cell extracts: Stomach 15%, Intestine 15%, esophagus 15%, pancreas 15%, embryonic endoderm 10%, Thymus 10%, Stabilizers and Mannitol 20%.</p> <p>Each 10-ml/250-mg Solvent Vial contains: Opotherapeutic cell extracts: Placenta 10%, 2% procaine, Sodium chloride 0.9%, Fructooligosaccharide 5%, enzyme complex: Superoxide dismutase, glutathione peroxidase, glutathione reductase, glutathione transferase, Stabilizers and sufficient sterilized Water for injection.</p>	<p>Formula components reach the cells directly or indirectly, in the case of oral products, by bloodstream, and are selectively incorporated into the cells through various means of cellular transport.</p> <p>It acts revitalizing the digestive system at the cellular level, improving its functionality and reducing the risk of degenerative diseases.</p> <div data-bbox="1387 1044 1713 1117" style="background-color: #800000; color: white; padding: 5px; text-align: center;"> Contraindications </div> <ul style="list-style-type: none"> • Allergies to animal proteins • Allergy to any of its components • Pregnancy and lactation 	<ul style="list-style-type: none"> • Mild colitis, gastritis, abdominal distension. <p>Orally: Two tablets in the morning and 2 at night, for at least six months.</p> <p>Intramuscular: 2 ml daily for 5 days, rest for two days and restart with 2 ml daily for 5 days. Repeat treatment at 3 months.</p> <ul style="list-style-type: none"> • Moderate colitis, constipation, malabsorption syndrome <p>Orally: Two tablets in the morning and 2 at night, for at least three months.</p> <p>Intramuscular: 2 ml daily for 5 days rest for two days and restart with 2 ml daily for 5 days. Repeat treatment with two more boxes.</p> <p>The tablets are taken in the morning on an empty stomach and at night before dinner (30 minutes before meals).</p> <p>NOTE: The dose may be increased according to the clinical picture of the patient and the physician's discretion; the results depend on the completion of treatment.</p>



Cellorgane Multicomplex® 3G

DIGESTIVE SYSTEM

Adjuvant treatment with:

Category	Therapeutic Class
Intestinal motility regulators	Cisapride, metoclopramide, Trimebutine, Pinaverium
Antacids	Omeprazole, famotidine, pantoprazole, ranitidine
Stool softeners	Fiber, Senosides, Docusate
Digestive enzymes	Bromelain, papain, pepsin, trypsin, lipase, Pancreatin